

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use HEPARIN SODIUM IN 0.45% SODIUM CHLORIDE INJECTION or HEPARIN SODIUM IN 5% DEXTROSE INJECTION safely and effectively. See full prescribing information for HEPARIN SODIUM IN 0.45% SODIUM CHLORIDE INJECTION or HEPARIN SODIUM IN 5% DEXTROSE INJECTION.

**HEPARIN SODIUM, for intravenous use**

Initial U.S. Approval: 1939

**Rx only****INDICATIONS AND USAGE**

Heparin sodium is an anticoagulant indicated for: (1)

- Prophylaxis and treatment of venous thromboembolism and pulmonary embolism
- Atrial fibrillation with embolization
- Treatment of acute and chronic consumptive coagulopathies (disseminated intravascular coagulation)
- Prevention of clotting in arterial and cardiac surgery
- Prophylaxis and treatment of peripheral arterial embolism
- Anticoagulant use in blood transfusions, extracorporeal circulation, and dialysis procedures.

**DOSAGE AND ADMINISTRATION**

Recommended Adult Dosages:

- Therapeutic Anticoagulant Effect with Full-Dose Heparin\* (2.3)

Intermittent Intravenous Injection	Initial Dose	10,000 units
	Every 4 to 6 hours	5,000 to 10,000 units
Continuous Intravenous Infusion	Initial Dose	5,000 units by intravenous injection
	Continuous	20,000 to 40,000 units/24 hours

\*Based on 150 lb. (68 kg) patient.

- Surgery of the Heart and Blood Vessels (2.5)

Intravascular via Total Body Perfusion	Initial Dose	≥ 150 units/kg; adjust for longer procedures
--	--------------	--

- Extracorporeal Dialysis (2.8)

Intravascular via Extracorporeal Dialysis	Follow equipment manufacturer’s operating directions carefully.
---	---

- See full prescribing information for recommended pediatric dosage. (2.4)

**DOSAGE FORMS AND STRENGTHS**

Heparin sodium is available as: (3)

***Heparin Sodium in 0.45% Sodium Chloride Injection:***

- Injection: 50 USP units per mL in 0.45% Sodium Chloride clear solution (25,000 USP units per 500 mL) in single-dose **freeflex**<sup>®</sup> bag
- Injection: 100 USP units per mL in 0.45% Sodium Chloride clear solution (25,000 USP units per 250 mL) in single-dose **freeflex**<sup>®</sup> bag

***Heparin Sodium in 5% Dextrose Injection:***

- Injection: 50 USP units per mL in 5% Dextrose clear solution (25,000 USP units per 500 mL) in single-dose **freeflex**<sup>®</sup> bag
- Injection: 100 USP units per mL in 5% Dextrose clear solution (25,000 USP units per 250 mL) in single-dose **freeflex**<sup>®</sup> bag

**CONTRAINDICATIONS**

- History of Heparin-Induced Thrombocytopenia (HIT) and Heparin-Induced Thrombocytopenia and Thrombosis (HITT) (4)
- Known hypersensitivity to heparin or pork products (4)
- In whom suitable blood coagulation tests cannot be performed at appropriate intervals (4)

**WARNINGS AND PRECAUTIONS**

- Fatal Medication Errors: Confirm choice of correct strength prior to administration. (5.1)
- Hemorrhage: Fatal cases have occurred. Use caution in conditions with increased risk of hemorrhage. (5.2)
- HIT and HITT: Monitor for signs and symptoms and discontinue if indicative of HIT and HITT. (5.3)
- Monitoring: Blood coagulation tests guide therapy for full-dose heparin. Monitor platelet count and hematocrit in all patients receiving heparin. (5.5)

**ADVERSE REACTIONS**

Most common adverse reactions are hemorrhage, thrombocytopenia, HIT and HITT, hypersensitivity reactions, and elevations of aminotransferase levels. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact**

**Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**DRUG INTERACTIONS**

Drugs that interfere with coagulation, platelet aggregation or drugs that counteract coagulation may induce bleeding. (7)

**See 17 for PATIENT COUNSELING INFORMATION.**

Revised: 12/2019

01-59-08-001B

**FULL PRESCRIBING INFORMATION: CONTENTS\*****1 INDICATIONS AND USAGE****2 DOSAGE AND ADMINISTRATION**

## 2.1 Preparation for Administration

## 2.2 Laboratory Monitoring for Efficacy and Safety

## 2.3 Therapeutic Anticoagulant Effect with Full-Dose Heparin

## 2.4 Pediatric Use

## 2.5 Cardiovascular Surgery

## 2.6 Converting to Warfarin

## 2.7 Converting to Oral Anticoagulants other than Warfarin

## 2.8 Extracorporeal Dialysis

**3 DOSAGE FORMS AND STRENGTHS****4 CONTRAINDICATIONS****5 WARNINGS AND PRECAUTIONS**

## 5.1 Fatal Medication Errors

## 5.2 Hemorrhage

## 5.3 Heparin-Induced Thrombocytopenia (HIT) and Heparin-Induced Thrombocytopenia and Thrombosis (HITT)

## 5.4 Thrombocytopenia

## 5.5 Coagulation Testing and Monitoring

## 5.6 Heparin Resistance

## 5.7 Hypersensitivity

**6 ADVERSE REACTIONS**

## 6.1 Postmarketing Experience

1593

**7 DRUG INTERACTIONS**

## 7.1 Oral Anticoagulants

## 7.2 Platelet Inhibitors

## 7.3 Other Interactions

**8 USE IN SPECIFIC POPULATIONS**

## 8.1 Pregnancy

## 8.2 Lactation

## 8.4 Pediatric Use

## 8.5 Geriatric Use

**10 OVERDOSAGE****11 DESCRIPTION****12 CLINICAL PHARMACOLOGY**

## 12.1 Mechanism of Action

## 12.2 Pharmacodynamics

## 12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

## 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**16 HOW SUPPLIED/STORAGE AND HANDLING****17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

- Use a non-vented infusion set or close the air-inlet on a vented set. The BLUE infusion port is compatible with spike systems produced according to ISO 8536-4, with an external spike diameter of 5.5 to 5.7 mm.
- Close the roller clamp of the infusion set.
- Hold the base of the BLUE infusion port and insert the spike by rotating your wrist slightly until the spike is fully inserted.
- The port membrane contains a self-sealing septum that helps prevent leakage after removing the spike. The infusion port is not intended to be spiked more than once.
- Hang from the hole at the top of the bag.
- For Single Use Only. Discard unused portion.

Do not admix with other drugs.

Do not use flexible container in series connections.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

**2.2 Laboratory Monitoring for Efficacy and Safety**

Adjust the dosage of heparin sodium according to the patient’s coagulation test results. When heparin is given by continuous intravenous infusion, determine the coagulation time approximately every 4 hours in the early stages of treatment. When the drug is administered intermittently by intravenous injection, perform coagulation tests before each injection during the early stages of treatment and at appropriate intervals thereafter. Dosage is considered adequate when the activated partial thromboplastin time (APTT) is 1.5 to 2 times the normal or when the whole blood clotting time is elevated approximately 2.5 to 3 times the control value.

Periodic platelet counts, hematocrits, and tests for occult blood in stool are recommended during the entire course of heparin therapy.

**2.3 Therapeutic Anticoagulant Effect with Full-Dose Heparin**

The dosing recommendations in Table 1 are based on clinical experience. Although dosage must be adjusted for the individual patient according to the results of suitable laboratory tests, the following dosage schedules may be used as guidelines:

**Table 1: Recommended Adult Full-Dose Heparin Regimens for Therapeutic Anticoagulant Effect**

Method of Administration	Frequency	Recommended Dose*
Intermittent Intravenous Injection	Initial Dose	10,000 units
	Every 4 to 6 hours	5,000 to 10,000 units
Continuous Intravenous Infusion	Initial Dose	5,000 units by intravenous injection
	Continuous	20,000 to 40,000 units per 24 hours

\* Based on 150 lb. (68 kg) patient.

**2.4 Pediatric Use**

There are no adequate and well-controlled studies on heparin use in pediatric patients. Pediatric dosing recommendations are based on clinical experience. In general, the following dosage schedule may be used as a guideline in pediatric patients:

Initial Dose:	75 to 100 units/kg (intravenous bolus over 10 minutes)
Maintenance Dose	Infants: 25 to 30 units/kg/hour; Infants < 2 months have the highest requirements (average 28 units/kg/hour) Children > 1 year of age: 18 to 20 units/kg/hour; Older children may require less heparin, similar to weight-adjusted adult dosage
Monitoring:	Adjust heparin to maintain aPTT of 60 to 85 seconds, assuming this reflects an anti-Factor Xa level of 0.35 to 0.70.

**2.5 Cardiovascular Surgery**

Patients undergoing total body perfusion for open-heart surgery should receive an initial dose of not less than 150 units of heparin sodium per kilogram of body weight. Frequently, a dose of 300 units per kilogram is used for procedures estimated to last less than 60 minutes or 400 units per kilogram for those estimated to last longer than 60 minutes.

**2.6 Converting to Warfarin**

To ensure continuous anticoagulation when converting from Heparin Sodium to warfarin, continue full heparin therapy for several days until the INR (prothrombin time) has reached a stable therapeutic range. Heparin therapy may then be discontinued without tapering [*see Drug Interactions (7.1)*].

**2.7 Converting to Oral Anticoagulants other than Warfarin**

For patients currently receiving intravenous heparin, stop intravenous infusion of heparin sodium immediately after administering the first dose of oral anticoagulant; or for intermittent intravenous administration of heparin sodium, start oral anticoagulant 0 to 2 hours before the time that the next dose of heparin was to have been administered.

**2.8 Extracorporeal Dialysis**

Follow equipment manufacturer’s operating directions carefully. A dose of 25 to 30 units/kg followed by an infusion rate of 1,500 to 2,000 units/hour is suggested based on pharmacodynamic data if specific manufacturers’ recommendations are not available.

**3 DOSAGE FORMS AND STRENGTHS**

Heparin Sodium in 0.45% Sodium Chloride Injection is available as:

- Injection: 50 USP units per mL in 0.45% Sodium Chloride clear solution (25,000 USP units per 500 mL) in single-dose **freeflex**<sup>®</sup> bag
- Injection: 100 USP units per mL in 0.45% Sodium Chloride clear solution (25,000 USP units per 250 mL) in single-dose **freeflex**<sup>®</sup> bag

Heparin Sodium in 5% Dextrose Injection is available as:

- Injection: 50 USP units per mL in 5% Dextrose clear solution (25,000 USP units per 500 mL) in single-dose **freeflex**<sup>®</sup> bag
- Injection: 100 USP units per mL in 5% Dextrose clear solution (25,000 USP units per 250 mL) in single-dose **freeflex**<sup>®</sup> bag

**4 CONTRAINDICATIONS**

The use of Heparin Sodium in 0.45% Sodium Chloride Injection or Heparin Sodium in 5% Dextrose Injection is contraindicated in patients with the following conditions:

- History of Heparin-Induced Thrombocytopenia (HIT) and Heparin-Induced Thrombocytopenia and Thrombosis (HITT) [*see Warnings and Precautions (5.3)*]
- Known hypersensitivity to heparin or pork products (e.g., anaphylactoid reactions) [*see Adverse Reactions (6.1)*]
- In whom suitable blood coagulation tests — e.g., the whole blood clotting time, partial thromboplastin time, etc., — cannot be performed at appropriate intervals (this contraindication refers to full-dose heparin; there is usually no need to monitor coagulation parameters in patients receiving low-dose heparin) [*see Warnings and Precautions (5.5)*]
- An uncontrolled bleeding state [*see Warnings and Precautions (5.2)*], except when this is due to disseminated intravascular coagulation.

**5 WARNINGS AND PRECAUTIONS****5.1 Fatal Medication Errors**

Do not use this product as a “catheter lock flush” product. Heparin is supplied in various strengths. Fatal hemorrhages have occurred due to medication errors. Carefully examine all heparin products to confirm the correct container choice prior to administration of the drug.

**5.2 Hemorrhage**

Avoid using heparin in the presence of major bleeding, except when the benefits of heparin therapy outweigh the potential risks.

Hemorrhage, including fatal events, has occurred in patients receiving Heparin Sodium. Hemorrhage can occur at virtually any site in patients receiving heparin. Adrenal hemorrhage (with resultant acute adrenal insufficiency), ovarian hemorrhage, and retroperitoneal hemorrhage have occurred during anticoagulant therapy with heparin [*see Adverse Reactions (6.1)*]. A higher incidence of bleeding has been reported in patients, particularly women, over 60 years of age [*see Clinical Pharmacology (12.3)*]. An unexplained fall in hematocrit or fall in blood pressure should lead to serious consideration of a hemorrhagic event.

Use heparin sodium with caution in disease states in which there is increased risk of hemorrhage, including:

- Cardiovascular** — Subacute bacterial endocarditis, severe hypertension.
- Surgical** — During and immediately following (a) spinal tap or spinal anesthesia or (b) major surgery, especially involving the brain, spinal cord or eye.
- Hematologic** — Conditions associated with increased bleeding tendencies, such as hemophilia, thrombocytopenia and some vascular purpuras.
- Patients with hereditary antithrombin III deficiency receiving concurrent antithrombin III therapy** — The anticoagulant effect of heparin is enhanced by concurrent treatment with antithrombin III (human) in patients with hereditary antithrombin III deficiency. To reduce the risk of bleeding, reduce the heparin dose during concomitant treatment with antithrombin III (human).
- Gastrointestinal** — Ulcerative lesions and continuous tube drainage of the stomach or small intestine.
- Other** — Menstruation, liver disease with impaired hemostasis.

